

# Viral Hepatitis

# Anatomy & physiology




☞ **Liver** is the largest gland in the body .

☞ **Located** behind the ribs in the right upper portion of the abdominal cavity .

☞ That manufactures, stores, alters, and excretes a large number of substances involved in metabolism.

☞ It removes the waste products from the blood stream and secretes them into the **Bile**.

Liver Anatomy demonstrate liver ducts and its relation with gallbladder ducts...



<b>Function</b>	<b>Actions</b>
<b>Digestion</b>	<ul style="list-style-type: none"> <li>➤ Bile salts for digestion/emulsification of fats</li> <li>➤ Processing and storage of fats, carbohydrates and proteins absorbed by the intestine and sent to the liver via the portal circulation</li> <li>➤ Processing and storage of vitamins and minerals</li> <li>➤ Vitamin B 12, A, D, E ,K</li> </ul>
<b>Endocrine</b>	<ul style="list-style-type: none"> <li>➤ Metabolism of glucocorticoids, mineralocorticoids and sex hormones</li> <li>➤ Regulation of fat, carbohydrate and protein metabolism</li> <li>➤ Glucose stored as glycogen</li> <li>➤ Main source of body heat</li> </ul>
<b>Hematologic</b>	<ul style="list-style-type: none"> <li>➤ Temporary storage of blood</li> <li>➤ Synthesis of bilirubin from breakdown of RBCs</li> <li>➤ Hematopoiesis in certain disease states</li> <li>➤ Synthesis of blood clotting factors (fibrinogen, thrombin, prothrombin)</li> <li>➤ synthesis of albumin, prealbumin,</li> </ul>
<b>Excretion</b>	<ul style="list-style-type: none"> <li>➤ Excretion of bile pigment</li> <li>➤ Excretion of cholesterol via bile</li> <li>➤ Urea synthesis as the final step in the excretion of ammonia (protein breakdown)</li> <li>➤ Detoxification of drugs, poisons and other foreign substances</li> </ul>

## **FUNCTIONS OF LIVER**



**Blood Detoxification  
& Purification**



**Glucose  
Regulation**





**Waste  
elimination**



**Fat  
Metabolism  
Breakdown**







is a systemic, viral infection in which necrosis and inflammation of liver cells produce a characteristic cluster of Clinical biochemical, and cellular changes.

# ➤ 5 Types for Viral Hepatitis

## Hepatitis

Inflammation of the Liver



Hepatitis

A

Hepatitis A Virus

Hepatitis

B

Hepatitis B Virus

Hepatitis

C

Hepatitis C Virus

Hepatitis

D

Hepatitis D Virus

Hepatitis

E

Hepatitis E Virus

<b>Virus Name</b>	<b>Incubation Period</b>	<b>Mode of transmission</b>	<b>Source of infection and spread of disease</b>
Hepatitis A virus ( HAV)	15-50 days ( average 30 days)	Fecal-oral route; poor sanitation, person to person contact. Waterborne; food borne.	Contaminated food, milk, water and selfish; person's with sub clinical infections infected food handler, poor personal hygiene, poor sanitation.
Hepatitis B virus ( HBV)	28-160 days (average 70-80 days)	<ul style="list-style-type: none"> <li>▪ Percutaneous (parenteral)/ per mucosal exposure to blood or blood products</li> <li>▪ Sexual contact.</li> <li>▪ Perinatal transmission</li> </ul>	<ul style="list-style-type: none"> <li>▪ Contaminated needles, syringes, and blood products.</li> <li>▪ Sexual activity with infected partners.</li> <li>▪ Oral-oral contact.</li> <li>▪ Tattoo / body piercing, bites.</li> <li>▪ Occupational hazards for health care personnel, hemodialysis staff, chemotherapy nurses, operating room nurses, dentists, persons at risk needle sticks.</li> </ul>
Hepatitis C virus ( HCV) (non A –non-B) Called post transfusion H.	15-160 days (average 50 days)	<ul style="list-style-type: none"> <li>▪ Transfusion of blood and blood products</li> <li>▪ Exposure to contaminated blood or blood products through equipment</li> <li>▪ High risk sexual contact.</li> <li>▪ Perinatal contact.</li> </ul>	<ul style="list-style-type: none"> <li>▪ Blood and blood products, needles and syringes.</li> <li>▪ Sexual activity with infected partners.</li> <li>▪ Increased with sexual transmitted disease.</li> <li>▪ Chronic treatment with hemodialysis.</li> </ul>
Hepatitis D virus ( HDV)	21-140 days (average 35days) <ul style="list-style-type: none"> <li>▪ HBV must precede HDV;</li> <li>▪ Chronic carriers of HBV are always at risk.</li> </ul>	<ul style="list-style-type: none"> <li>▪ Can cause infection only together with HBV.</li> <li>▪ Rotes of transmission same as for HBV.</li> <li>▪ HBV surface antigen necessary for replication.</li> </ul>	<ul style="list-style-type: none"> <li>▪ Same as HBV.</li> </ul>
Hepatitis E virus ( HEV) Similar to HAV	15-65days ( average 42 days)	<ul style="list-style-type: none"> <li>▪ Fecal-oral.</li> <li>▪ Outbreaks associated with contaminated water supply in developing countries</li> </ul>	<ul style="list-style-type: none"> <li>▪ Contaminated water; poor sanitation</li> </ul>

## **Clinical Manifestation of the Phases of Hepatitis:**

<b><u>Preicteric</u></b> <b><u>It lasts 1-21 days</u></b>	<b><u>Icteric</u></b> <b><u>It lasts 2-4 weeks</u></b>	<b><u>Post Icteric</u></b> <b><u>It persists several months</u></b>
<ol style="list-style-type: none"><li>1. Anorexia.</li><li>2. Nausea, vomiting.</li><li>3. Right upper quadrant discomfort.</li><li>4. Decrease sense of taste and smell.</li><li>5. Malaise.</li><li>6. Headache.</li><li>7. Fever.</li><li>8. Arthralgias.</li><li>9. Urticaria, rash.</li><li>10. Hepatomegaly &amp; tenderness of the liver.</li><li>11. Splenomegaly.</li><li>12. Weight loss.</li></ol>	<ol style="list-style-type: none"><li>1. Jaundice.</li><li>2. Pruritus due to accumulation of bile salts under skin.</li><li>3. Dark urine.</li><li>4. Bilirubinuria.</li><li>5. Light or clay colored stool.</li><li>6. Fatigue.</li><li>7. Continued hepatomegaly with tenderness.</li><li>8. Weight loss.</li></ol>	<ol style="list-style-type: none"><li>1- Malaise.</li><li>2- Easy fatigability.</li><li>3- Hepatomegaly.</li></ol>

---

# Past health history:



- ❧ Previous liver disease, hepatitis immunization.
- ❧ Hemophilia.(An inherited bleeding disease)
- ❧ Exposure to infected persons.
- ❧ Ingestion of contaminated food and water.
- ❧ Sexual promiscuity.
- ❧ Exposure to toxins.
- ❧ Exposure to contaminated needles.
- ❧ Recent travel.
- ❧ Organ transplant recipient.
- ❧ Exposure to new drug regimen.

- **Present History**



**Fatigue.**

**Weight loss.**

**Digestive  
disturbance.**

**Skin changes.**

**Feeling of  
fullness in  
right upper  
quadrant.**

# • Objective Data:



☞ Vital signs: Hypertension, tachypnea, low grade fever.

☞ Skin: Dryness, scratches, jaundice, bruises

Angioedema :(Sever form of urticaria which involve skin, face, hands and genital organ)

☞ Eyes: Icteric sclera.

☞ Thorax: Spider angiomas.

☞ Abdomen: Distention, prominent veins, hepatomegaly, Splenomegaly.

# • Possible findings(Diagnostic tests):

- ❧ Abnormal liver enzyme studies.
- ❧ Elevated serum bilirubin.
- ❧ Hypoalbuminemia.
- ❧ Anemia.
- ❧ Bilirubin in urine and increased urobilinogen.
- ❧ Prolonged prothrombin time.
- ❧ Serologic test positive for hepatitis, including anti-HAV IgM, anti-HBc IgM, anti-HCV, anti-HDV.
- ❧ Abnormal liver scan.
- ❧ Positive liver biopsy.

## Preventive Measures of Viral Hepatitis

### **Hepatitis A**

#### General Measures:

- Hand washing.
- Proper personal hygiene.
- Environmental sanitation.
- Safe practice for preparing and dispensing food
- Control and screening (signs and symptoms of food handlers).
- Serologic screening while carrying virus.
- Active immunization.
- Community health education program.

#### Use of immune globulin:

- Early administration (1-2 weeks after exposure to those exposed).
- Use of prophylaxis for travelers to areas where hepatitis A is common.

### **Hepatitis B and C**

#### Percutaneous Transmission:

- Screening of donated blood.
- Use of disposable needles and syringes

#### Sexual Transmission:

- Acute exposure: HBIG administration to sexual partner of HbsAg-positive person.
- Use condom for sexual intercourse.

#### General Measures:

- Hand washing.
- Avoid sharing tooth brushes and razors
- HBIG administration for one-time exposure (needle stick, contact of mucous membranes with infectious material).
- Active immunization: HBV vaccine.

# What is the most common treatment for viral hepatitis?

---

There is no cure for hepatitis B, which resolves on its own in 95% of cases. Supportive care can help manage symptoms. In cases of chronic illness, a doctor may prescribe an **antiviral medication**, and they will monitor the liver regularly to check for damage over time.

**Measures to Prevent Transmission of Hepatitis Viruses From Patients to health Care Personnel:**


<b>Hepatitis A</b>	<b>Hepatitis B</b>	<b>Hepatitis C</b>
<ul style="list-style-type: none"><li>▪ Always maintain good personal hygiene.</li><li>▪ Wash hands after contact with a patient or removal of gloves.</li><li>▪ Use infection control precautions.</li></ul>	<ul style="list-style-type: none"><li>▪ Use infection control precautions.</li><li>▪ Wash hands.</li><li>▪ Reduce contact with blood or blood-containing secretions.</li><li>▪ Handle the blood of patients as potentially infective.</li><li>▪ Dispose of needles properly.</li><li>▪ Administer HBV vaccine to all health care personnel.</li><li>▪ Use needles IV access devices when available.</li></ul>	<ul style="list-style-type: none"><li>▪ Use infection control precaution.</li><li>▪ Wash hands.</li><li>▪ Reduce contact with blood or blood contaminated secretions.</li><li>▪ Handle the blood of patient as potentially infective.</li><li>▪ Dispose needles properly.</li><li>▪ Use needlesless IV access devices when available.</li></ul>





## **Nursing Diagnosis**

- Altered Nutrition: Less Than Body Requirements related to anorexia as manifested by inadequate food intake.
- Fluid Volume Deficit related to nausea and/or vomiting



# Hepatitis

**H**andwashing (**strict**)

**e**at **LOW** fat & **HIGH** carbs

**p**ersonal hygiene products **NOT** shared

**a**ctivity conservation (**REST!!**)

**t**oxic substances **avoided**  
-**alcohol**, **sedatives**, aspirin, **acetaminophen** etc.

**i**ndividual bathroom

**t**esting results

**i**nterferon (**subcutaneous**)

**S**mall but frequent meals

